California Association for Bilingual Education TB Compliance Form		
Name:		Date of Birth:
Job Title:		
Social Security No:		Email Address:
APPLICANTS : Date of x-	-ray must be within six months prisk factors, or if risk factors	a negative chest x-ray on(date). brior to date of hire. were identified, the patient has been examined and
Health Care Provider Signature (MD, DO, PA, NP, RN	<u>ONLY</u>)	Date
Print Health Care Provider's Name	Title	License No.
Address:	City	Zip Code
	Fax	
Telephone	1 444	

Adult Tuberculosis (TB) Risk Assessment Questionnaire

(To satisfy California Education Code Section 49406 and Health and Safety Code Sections 121525-121555) To be administered by a licensed health care provider (physician, physician assistant, nurse practitioner, registered nurse)

- The purpose of this tool is to identify <u>adults</u> with infectious tuberculosis (TB) to prevent them from spreading disease.
- Do not repeat testing unless there are <u>new risk factors since the last negative test</u>.
- Do not treat for latent TB infection (LTBI) until active TB disease has been excluded:

For individuals with signs or symptoms of TB disease or abnormal chest x-ray consistent with TB disease, evaluate for active TB disease with a chest x-ray, symptom screen, and if indicated, sputum AFB smears, cultures and nucleic acid amplification testing.

A negative tuberculin skin test (TST) or interferon gamma release assay (IGRA) does not rule out active TB disease.

Name of Person Assessed for TB Risk Factors: ______

Assessment Date: _____

Date of Birth: _____

History of Tuberculosis Disease or Infection (Check appropriate box below)

Yes

• If there is a documented history of positive TB test or TB disease, then a symptom review and chest x-ray (if none performed in

the previous 6 months) should be performed at initial hire by a physician, physician assistant, or nurse practitioner. If the x-ray does not have evidence of TB, the person is no longer required to submit to a TB risk assessment or repeat chest x-rays.

No (Assess for Risk Factors for Tuberculosis using box below)

TB testing is recommended if any of the 3 boxes below are checked

One or more sign(s) or symptom(s) of TB disease

- TB symptoms include prolonged cough, coughing up blood, fever, night sweats, weight loss, or excessive fatigue.
- **Birth, travel, or residence** in a country with an elevated TB rate for at least 1 month
 - Includes countries other than the United States, Canada, Australia, New Zealand, or Western and North European countries.
 - Interferon gamma release assay (IGRA) is preferred over tuberculin skin test (TST) for non-US-born persons.

Close contact to someone with infectious TB disease during lifetime

Treat for LTBI if TB test result is positive and active TB disease is ruled out

Do Not Submit Adult Tuberculosis (TB) Risk Assessment Questionnaire to CABE